

OTUB2 Protein, Human (His)

Cat. No.:	HY-P77118
Synonyms:	Ubiquitin thioesterase OTUB2; Otubain-2; C14orf137; OTB2; OTU2
Species:	Human
Source:	E. coli
Accession:	Q96DC9 (M1-H234)
Gene ID:	78990
Molecular Weight:	Approximately 28 kDa

PROPERTIES

AA Sequence	<pre> M S E T S F N L I S E K C D I L S I L R D H P E N R I Y R R K I E E L S K R F T A I R K T K G D G N C F Y R A L G Y S Y L E S L L G K S R E I F K F K E R V L Q T P N D L L A A G F E E H K F R N F F N A F Y S V V E L V E K D G S V S S L L K V F N D Q S A S D H I V Q F L R L L T S A F I R N R A D F F R H F I D E E M D I K D F C T H E V E P M A T E C D H I Q I T A L S Q A L S I A L Q V E Y V D E M D T A L N H H V F P E A A T P S V Y L L Y K T S H Y N I L Y A A D K H </pre>
Biological Activity	The specific activity was determined to be 73.477 nmol/min/mg in a DUB assay.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 50 mM Tris-HCL, 300 mM NaCl, 500 mM arginine, pH 10.0.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	As a hydrolase, OTUB2 exhibits the capacity to efficiently remove conjugated ubiquitin from proteins in vitro, suggesting a pivotal regulatory role in protein turnover by preventing degradation. With its deubiquitination activity, OTUB2 acts on 'Lys-11', 'Lys-48', and 'Lys-63'-linked polyubiquitin chains, displaying a preference for the latter. This specificity underscores OTUB2's selectivity in modulating diverse ubiquitin linkages, emphasizing its potential significance in orchestrating cellular
-------------------	--

processes related to protein ubiquitination and degradation.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA