**Proteins** 





## Prostasin/PRSS8 Protein, Rat (HEK293, His)

Cat. No.: HY-P77155

Synonyms: Channel-activating protease 1; CAP1; Serine protease 8

Species: Rat

HEK293 Source:

Q9ES87 (A30-R322) Accession:

Gene ID: 192107

Molecular Weight: Approximately 32.9 kDa.

## **PROPERTIES**

AA Sequence				
·	ADGTEASCGA	VIQPRITGGG	SAKPGQWPWQ	$V \; S \; I \; T \; Y \; N \; G \; V \; H \; V$
	$C\;G\;G\;S\;L\;V\;S\;N\;Q\;W$	VVSAAHCFPR	EHSKEEYEVK	LGAHQLDSFS
	NDIVVHTVAQ	IISHSSYREE	GSQGDIALIR	LSSPVTFSRY
	IRPICLPAAN	ASFPNGLHCT	VTGWGHVAPS	VSLQTPRPLQ
	QLEVPLISRE	TCSCLYNINA	VPEEPHTIQQ	DMLCAGYVKG
	GKDACQGDSG	GPLSCPIDGL	WYLAGIVSWG	DACGAPNRPG
	VYTLTSTYAS	WIHHHVAELQ	PRVVPQTQES	QPDGHLCNHH

PVFNLAAAQK LSR

**Biological Activity** Measured by its ability to cleave the fluorogenic peptide substrate Boc-QAR-AMC. The specific activity is 31.804 pmoL/min/µg, as measured under the described conditions.

Lyophilized powder **Appearance** 

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

**Endotoxin Level** <1 EU/ $\mu$ g, determined by LAL method.

Reconsititution It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH<sub>2</sub>O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).

Storage & Stability Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.

**Shipping** Room temperature in continental US; may vary elsewhere.

## **DESCRIPTION**

**Background** Prostasin/PRSS8, characterized by a trypsin-like cleavage specificity with a preference for poly-basic substrates, plays a

Page 1 of 2

crucial role in stimulating the activity of the epithelial sodium channel (ENaC). Its activation involves the cleavage of the gamma subunits (SCNN1G) to modulate ion transport processes. Structurally, Prostasin/PRSS8 forms a heterodimer comprising light and heavy chains, united by a disulfide bond, underscoring its functional complexity in mediating cellular responses.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com