

Proteins

Screening Libraries

Inhibitors

Product Data Sheet

Nectin-1 Protein, Rat (HEK293, Fc)

Cat. No.: HY-P77470

Poliovirus Receptor-Related Protein 1; Herpes Virus Entry Mediator C; Herpesvirus Entry Synonyms:

Mediator C; HveC; Herpesvirus Ig-Like Receptor; HlgR; Nectin-1; CD111; PVRL1; HVEC; PRR1

Species:

HEK293 Source:

Accession: NP_001093946 (M1-A354)

Gene ID: 192183 Molecular Weight: 88-98 kDa

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu g/mL$ in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

Nectin-1 Protein, characterized by its carbohydrate binding and virion binding activities, plays pivotal roles in axon guidance, regulation of synapse assembly, and viral entry into host cells. Distributed across various cellular components, including dendrites, growth cone membranes, and synapses, Nectin-1 is implicated in human conditions such as cleft lip, cleft lip-palate-ectodermal dysplasia syndrome, cleft palate, and ectodermal dysplasia. The biased expression pattern, notably in the Kidney (RPKM 45.4), Brain (RPKM 31.3), and nine other tissues, underscores its significance in diverse physiological contexts.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 1 of 1