

DR3/TNFRSF25 Protein, Human (Biotinylated, HEK293, Fc)

| | |
|--------------------------|---|
| Cat. No.: | HY-P77522 |
| Synonyms: | Tumor necrosis factor receptor superfamily member 25; Apo-3; LARD; TNFRSF25; DR3; TNFRSF12; WSL; WSL1 |
| Species: | Human |
| Source: | HEK293 |
| Accession: | Q93038 (Q25-Q199) |
| Gene ID: | 8718 |
| Molecular Weight: | Approximately 45.9 kDa. |

PROPERTIES

| | |
|--------------------------------|---|
| AA Sequence | <p> Q G G T R S P R C D C A G D F H K K I G L F C C R G C P A G H Y L K A P C T E P C G N S T C L V C P Q D T F L A W E N H H N S E C A R C Q A C D E Q A S Q V A L E N C S A V A D T R C G C K P G W F V E C Q V S Q C V S S S P F Y C Q P C L D C G A L H R H T R L L C S R R D T D C G T C L P G F Y E H G D G C V S C P T S T L G S C P E R C A A V C G W R Q </p> |
| Appearance | Lyophilized powder |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. |
| Endotoxin Level | <1 EU/µg, determined by LAL method. |
| Reconstitution | It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage. |
| Shipping | Room temperature in continental US; may vary elsewhere. |

DESCRIPTION

| | |
|-------------------|--|
| Background | <p>DR3/TNFRSF25 Protein serves as the receptor for TNFSF12/APO3L/TWEAK and directly interacts with the adapter TRADD. This interaction leads to the activation of NF-kappa-B and induction of apoptosis. The protein may play a crucial role in regulating lymphocyte homeostasis. It forms homodimers and exhibits strong interactions via the death domains with TNFRSF1 and TRADD, initiating distinct signaling cascades involved in apoptosis and NF-kappa-B signaling. Additionally, DR3/TNFRSF25 interacts with BAG4, contributing to its multifaceted roles in cellular responses.</p> |
|-------------------|--|

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA