

GDF-15 Protein, Cynomolgus (Biotinylated, HEK293, Fc)

Cat. No.:	HY-P77682
Synonyms:	GDF-15; MIC-1; NAG-1; PDF; PLAB; PTGFB; GDF15; MIC1; RG-1; Placental TGF-beta; PTGF-beta; PTGFBPTGF-beta
Species:	Cynomolgus
Source:	HEK293
Accession:	G7PWZ3 (A197-V308)
Gene ID:	102117226
Molecular Weight:	40-50 kDa

PROPERTIES

Biological Activity	Immobilized Cynomolgus GFRAL, His Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Cynomolgus GDF15, hFc Tag with the EC ₅₀ of 4.8ng/ml determined by ELISA.
Appearance	Solution.
Formulation	Supplied as a 0.22 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	N/A.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice.

DESCRIPTION

Background

Growth differentiation factor 15 (GDF-15) is a polypeptide hormone belonging to the transforming growth factor β (TGF-β) superfamily. GDF-15 was highly expressed in placenta, low in prostate and colon, and to some extent in kidney. So GDF-15 is also known as placental transforming growth factor PGF-β, placental bone morphogenetic protein PLAB, and prostatic-derived factor PDF. GDF-15 has a wide range of biological functions in physiology and pathology, especially in aging, cancer, and metabolic processes. GDF-15 is initially stored in the extracellular matrix (ECM), where it undergoes proteolytic hydrolysis upon external stimulation to form an active form that is quickly secreted into circulation. In mouse cardiomyocytes, the cleavage process of GDF-15 may be catalyzed by the enzymes of the PCSK family, resulting in a mature dimer form. Upstream of the GDF15 promoter site, there are binding sites for various transcription factors, including specific protein 1 (Sp1), early growth response protein 1 (Egr-1), p53 and COUP transcription factor 1 (COUP-TF1). The receptor of GDF-15 is alpha-like protein (GFRAL), a receptor of the glial cell derived neurotrophic factor (GDNF) family. The GFRAL-GDF15 complex binds to the tyrosine kinase co-receptor RET, leading to RET phosphorylation. Subsequently, GFRAL-GDF15 continued to activate the intracellular signaling pathways of AKT, ERK1/2, and phospholipase C (PLCγ), but not the SMAD pathway. GDF-15 is overexpressed during and after many pathological states such as tissue injury and inflammation. The

stimulating factors that contribute to this result include oxidized low-density lipoprotein (oxLDL), cytokines, and growth factors such as IL-1 β , TNF- α , angiotensin II, macrophage colony-stimulating factor M-CSF, and TGF β . GDF-15, also known as the NSAIDS drug activator gene NAG-1, may play an anti-inflammatory role by inhibiting macrophage activation. GDF-15 also inhibits the activity of NF κ B or the expression of several cytokines, including interferon (IFN- γ), interleukin-6 (IL-6), monocyte chemoattractant protein-1 (MCP-1), and tumor necrosis factor- α (TNF- α). GDF-15 has significant resistance to endotoxin-induced sepsis caused by acute kidney injury (AKI) and myocardial dysfunction. GDF-15 also appears to promote tumor growth in the later stages of malignancy. Elevated serum GDF15 levels have been reported as potential biomarkers for cancer progression, including breast, colon, pancreatic, and prostate tumors, among others. In human and cynomolgus monkeys, the amino acid sequence similarity of GDF-15 protein was high, and the similarity rate was 91.56%. Compared with the amino acid sequences of mice and rats, the similarity of human GDF-15 was low (59.73% and 59.39%, respectively) [1].

REFERENCES

[1]. Assadi A, et al. GDF15, an update of the physiological and pathological roles it plays: a review. *Pflügers Arch.* 2020 Nov;472(11):1535-1546.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA