

HLA-G Complex Tetramer Protein, Human (Biotinylated, HEK293, His-Avi)

Cat. No.:	HY-P77691
Synonyms:	HLA G antigen; sHLA-G; b2 microglobulin; HLA G; HLAG; HLA-G; MHC Class I Antigen G; MHC class Ib antigen; MHC-G; sHLA-G
Species:	Human
Source:	HEK293
Accession:	P17693 (G25-T305)&P61769 (I21-M119)&RIIPRHLQL
Gene ID:	3135&567
Molecular Weight:	260-265 kDa

PROPERTIES

Biological Activity	<ol style="list-style-type: none"> Human LILRB2 captured on CM5 Chip via Protein A can bind Biotinylated Human HLA-G Complex Tetramer with an affinity constant of 4.38 nM as determined in SPR assay (Biacore T200). Serial dilutions of Anti-LILRB2 Antibody were added into Biotinylated HLA-G Complex Tetramer, His Tag: Human LILRB2, mFc Tag binding reactions. The half maximal inhibitory concentration (IC₅₀) is 0.3 µg/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>HLA-G, a non-classical major histocompatibility class Ib molecule, plays a crucial role in immune regulation at the maternal-fetal interface. In association with B2M/beta-2 microglobulin, it forms a complex that selectively binds a limited repertoire of nonamer self-peptides derived from intracellular proteins, including histones and ribosomal proteins. This peptide-bound HLA-G-B2M complex acts as a ligand for inhibitory/activating KIR2DL4, LILRB1, and LILRB2 receptors on uterine immune cells, fostering fetal development while maintaining maternal-fetal tolerance. Interactions with KIR2DL4 and LILRB1 receptors trigger NK cell senescence-associated secretory phenotype, promoting vascular remodeling and fetal growth during early pregnancy. Moreover, HLA-G's engagement with LILRB2 induces the differentiation of type 1 regulatory T cells and myeloid-derived suppressor cells, actively contributing to the maintenance of maternal-fetal tolerance. Additionally, HLA-G may play a role in balancing tolerance and antiviral immunity by modulating the effector functions of NK cells, CD8+ T cells, and B cells. Furthermore, it negatively regulates NK cell- and CD8+ T cell-mediated cytotoxicity, highlighting its</p>
-------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

multifaceted role in immune regulation at the maternal-fetal interface.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA