

## IL-17A-17F Heterodimer Protein, Human (Biotinylated, HEK293, His-Avi)

Cat. No.:	HY-P77713
Synonyms:	IL-17A; Interleukin-17A; CTLA-8; IL-17; IL-17F; IL24; ML-1; Interleukin-17F
Species:	Human
Source:	HEK 293
Accession:	Q16552 (G24-A155)&Q96PD4 (R31-Q163)
Gene ID:	3605&112744
Molecular Weight:	Approximately 16 kDa

### PROPERTIES

Biological Activity	Immobilized Biotinylated Human IL-17A&F at 1 µg/mL (100µL/Well) on the plate. Dose response curve for Human IL-17R alpha with the EC <sub>50</sub> of 0.19 µg/mL determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA