

Product Data Sheet

SIRP alpha V3 Protein, Human (Biotinylated, H54L, HEK293, His-Avi)

Cat. No.:	HY-P77842
Synonyms:	CD172a; BIT; MFR; MYD1; MYD-1; P84; PTPNS1; SHP substrate 1; SHPS1; SHPS-1; SHPS1CD172A; SIRP alpha; SIRPA; Sirp-alpha-1; SIRPalpha2; Sirp-alpha-2; Sirp-alpha-3
Species:	Human
Source:	HEK293
Accession:	AAH38510 (E31-R369, H54L)
Gene ID:	140885
Molecular Weight:	55-65 kDa

PROPERTIESBiological ActivityImmobilized Biotinylated Human SIRP alpha V3, His Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Human CD47, hFc Tag with the EC ₅₀ of 0.12µg/ml determined by ELISA.AppearanceLyophilized powder.FormulationLyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.Endotoxin Level<1 EU/µg, determined by LAL method.ReconsititutionIt is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH20.Storage & StabilityStored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.ShippingRoom temperature in continental US; may vary elsewhere.	PROPERTIES	
CD47, hFc Tag with the EC ₅₀ of 0.12µg/ml determined by ELISA. Appearance Lyophilized powder. Formulation Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization. Endotoxin Level <1 EU/µg, determined by LAL method. Reconsititution It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. Storage & Stability Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.	PROPERTIES	
Formulation Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization. Endotoxin Level <1 EU/µg, determined by LAL method.	Biological Activity	
is of primitical nervel b) oprimitical nervel of Logic and Dial primitical contraction of Dial primitical co	Appearance	Lyophilized powder.
Reconsititution It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. Storage & Stability Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.	Formulation	
Storage & Stability Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.	Endotoxin Level	<1 EU/µg, determined by LAL method.
aliquots at -20°C or -80°C for extended storage.	Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu\text{g}/\text{mL}$ in ddH_2O.
Shipping Room temperature in continental US; may vary elsewhere.	Storage & Stability	
	Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA