Proteins



Product Data Sheet

ENPP-1 Protein, Human (HEK293, His)

Cat. No.: HY-P77923

Synonyms: E-NPP 1; NPPase; ENPP1; NPPS; PC1; PDNP1; Ly-41 Antigen; M6S1; M6S1NPP1; NPP1; PCA1;

Species: Human Source: HEK293

Accession: P22413 (K98-D925)

Gene ID: 5167

Molecular Weight: 110-120 kDa

PROPERTIES

AA Sequence	KPSCAKEVKS	CKGRCFERTF	GNCRCDAACV	ELGNCCLDYO
	ETCIEPEHIW	TCNKFRCGEK	RLTRSLCACS	DDCKDKGDCC
	INYSSVCQGE	KSWVEEPCES	INEPQCPAGF	ETPPTLLFSL
	DGFRAEYLHT	WGGLLPVISK	LKKCGTYTKN	MRPVYPTKTF
	PNHYSIVTGL	YPESHGIIDN	KMYDPKMNAS	FSLKSKEKFN
	PEWYKGEPIW	VTAKYQGLKS	GTFFWPGSDV	EINGIFPDIY
	KMYNGSVPFE	ERILAVLOWL	OLPKDERPHF	YTLYLEEPDS
	SGHSYGPVSS	EVIKALQRVD	GMVGMLMDGL	KELNLHRCLN
	LILISDHGME	QGSCKKYIYL	NKYLGDVKNI	KVIYGPAARL
	RPSDVPDKYY	SFNYEGIARN	LSCREPNQHF	KPYLKHFLPK
	RLHFAKSDRI	EPLTFYLDPQ	WQLALNPSER	KYCGSGFHGS
	DNVFSNMQAL	FVGYGPGFKH	GIEADTFENI	EVYNLMCDLL
	NLTPAPNNGT	HGSLNHLLKN	PVYTPKHPKE	VHPLVQCPFT
	RNPRDNLGCS	CNPSILPIED	FQTQFNLTVA	EEKIIKHETL
	PYGRPRVLQK	ENTICLLSQH	QFMSGYSQDI	LMPLWTSYTV
	DRNDSFSTED	FSNCLYQDFR	IPLSPVHKCS	FYKNNTKVSY
	GFLSPPOLNK	NSSGIYSEAL	LTTNIVPMYO	SFQVIWRYFH
	DTLLRKYAEE	RNGVNVVSGP	VFDFDYDGRC	DSLENLRQKR
	RVIRNQEILI	PTHFFIVLTS	CKDTSQTPLH	CENLDTLAFI
	LPHRTDNSES	CVHGKHDSSW	VEELLMLHRA	RITDVEHITG
	LSFYQQRKEP	VSDILKLKTH	LPTFSQED	RIIDVLIIIIG
	LSTIQQKKLT	VSDIERERIII		
Biological Activity	Measured by its ability to hydrolyze thymidine 5'-monophosphate p-nitrophenyl ester. The specific activity is 117007.4074			ester. The specific activity is 117007.4074
	pmol/min/μg, as measure	ed under the described cond	itions.	
Appearance	Lyophilized powder.			
Formulation	Lyophilized from 0.22 μ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.			
Endotoxin Level	<1 EU/µg, determined by LAL method.			

Page 1 of 2

Reconsititution	It is not recommended to reconstitute to a concentration less than 100 μ g/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

ENPP-1 Protein, a nucleotide pyrophosphatase, plays a crucial role in bone mineralization and soft tissue calcification by regulating pyrophosphate levels. By preferentially hydrolyzing ATP and other nucleoside 5' triphosphates, such as GTP, CTP, and UTP, ENPP-1 generates diphosphate (PPi), which inhibits the further growth of nascent hydroxyapatite crystals, thereby preventing excessive bone mineralization and calcification of soft tissues. Additionally, it participates in the hydrolysis of diadenosine polyphosphates and 3',5'-cAMP to AMP, and may regulate nucleotide sugar availability in the endoplasmic reticulum and Golgi, influencing purinergic signaling. ENPP-1 also contributes to the repression of hedgehog signaling, inhibiting ectopic joint calcification and maintaining articular chondrocytes. Furthermore, it is implicated in modulating insulin sensitivity and function, participating in melanogenesis, and can hydrolyze 2',3'-cGAMP, a second messenger involved in triggering type-I interferon production. The exact role of 2',3'-cGAMP hydrolysis in the extracellular space remains unclear.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA