

CD276/B7-H3 Protein, Human (Biotinylated, HEK293, His-Avi)

Cat. No.:	HY-P78074
Synonyms:	B7-H3; CD276; PSEC0249; 4Ig-B7-H3; UNQ309; PRO352; B7 homolog 3; CD276
Species:	Human
Source:	HEK293
Accession:	Q5ZPR3-1 (G27-T461)
Gene ID:	80381
Molecular Weight:	70-80 kDa

PROPERTIES

Biological Activity	<ol style="list-style-type: none"> 1. Immobilized Biotinylated Human B7-H3 (4Ig) His at 1 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-B7-H3 Ab hFc with the EC₅₀ ≤ 15.2 ng/mL determined by ELISA. 2. Immobilized Anti-B7-H3 Antibody, hFc Tag at 0.5 µg/mL (100 µl/well) on the plate. Dose response curve for Biotinylated Human B7-H3 (4Ig) , His Tag with the EC₅₀ ≤ 9.7 ng/mL determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years from date of receipt. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>The CD276/B7-H3 protein is suggested to play a multifaceted role in the regulation of T-cell-mediated immune responses, potentially acting as a protective factor in tumor cells by inhibiting natural-killer-mediated cell lysis and serving as a marker for the detection of neuroblastoma cells. Additionally, CD276/B7-H3 may be involved in the development of acute and chronic transplant rejection, contributing to the regulation of lymphocytic activity at mucosal surfaces. Notably, it could play a crucial role in providing the placenta and fetus with an immunologically suitable environment throughout pregnancy. Both isoform 1 and isoform 2 of CD276/B7-H3 appear redundant in their ability to modulate CD4 T-cell responses, with isoform 2 demonstrated to enhance the induction of cytotoxic T-cells and selectively stimulate interferon-gamma production in the presence of T-cell receptor signaling. The interaction with TREML2 is identified as enhancing T-cell activation, highlighting the diverse roles CD276/B7-H3 may play in immune regulation and cellular responses.</p>
-------------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA