

Certificate of Analysis

IL-17A Protein, Human (Biotinylated, 132a.a, HEK293, His-Avi)

| Cat. No.: | HY-P78157 |
|-------------------|-------------------|
| Batch No.: | 317967 |
| Species: | Human |
| Source: | HEK293 |
| Tag: | C-Avi;C-His |
| Accession: | Q16552 (G24-A155) |
| Gene ID: | 3605 |
| Molecular Weight: | 16-30 kDa |

ANALYTICAL DATA

| TEST | Specifications | Results |
|---------------------|--|------------|
| Purity | Greater than 95% as determined by Tris-Bis PAGE. | ≥95.0% |
| Endotoxin Level | <1 EU/µg, determined by LAL method. | PASS |
| Biological Activity | Immobilized Biotinylated Human IL-17A, His Tag at 2 µg/mL (100 µl/well) on the streptavidin precoated plate (5 µg/ml). Dose response curve for Human IL- 17R alpha, hFc Tag with the EC ₅₀ of ≤27.0 ng/mL determined by ELISA. | 15.1 ng/mL |
| Appearance | Lyophilized powder | |
| Reconstitution | It is not recommended to reconstitute to a concentration less than 100 $\mu\text{g}/\text{mL}$ in ddH_2O. | |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or - 20°C for longer (with carrier protein). It is recommended to freeze aliquots at - 20°C or -80°C for extended storage. | |
| Shipping | Room temperature in continental US; may vary elsewhere. | |

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

28-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA