

## Latent TGF beta 1/TGFB1 Protein, Human (Biotinylated, HEK293, His-Avi)

Cat. No.:	HY-P78214
Synonyms:	CEDLAP; DPD1; TGF beta1; TGFB; TGFB1; TGFBeta; TGF-beta-1; TGF β1; TGFβ; TGF-β-1
Species:	Human
Source:	HEK293
Accession:	P01137 (L30-S390, C33S)
Gene ID:	7040
Molecular Weight:	40-50&13-15 kDa

### PROPERTIES

Biological Activity	Immobilized Human ITGAV&ITGB6 His at 1 µg/mL (100µL/Well) on the plate. Dose response curve for Biotinylated Human Latent TGF beta 1 His with the EC <sub>50</sub> of ≤ 0.6 µg/mL determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	Latent TGF beta 1/TGFB1 Protein promotes fibrillin-1 and -2 assembly downstream of fibronectin in mouse embryonic fibroblasts <sup>[3]</sup> .
------------	------------------------------------------------------------------------------------------------------------------------------------------------

### REFERENCES

- [1]. J Taipale, et al. Latent transforming growth factor-beta 1 associates to fibroblast extracellular matrix via latent TGF-beta binding protein. J Cell Biol. 1994 Jan;124(1-2):171-81.
- [2]. C Unsöld, et al. Latent TGF-beta binding protein LTBP-1 contains three potential extracellular matrix interacting domains. J Cell Sci. 2001 Jan;114(Pt 1):187-197.
- [3]. Matthias Przyklenk, et al. LTBP1 promotes fibrillin incorporation into the extracellular matrix. bioRxiv. 2022.

---

[4]. Rui Cai, et al. LTBP1 promotes esophageal squamous cell carcinoma progression through epithelial-mesenchymal transition and cancer-associated fibroblasts transformation. J Transl Med. 2020 Mar 26;18(1):139.

[5]. Xiaojun Fu, et al. LTBP1 plays a potential bridge between depressive disorder and glioblastoma. J Transl Med. 2020 Oct 15;18(1):391.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA