

TRAIL R2/TNFRSF10B Protein, Human (Biotinylated, HEK293, His-Avi)

Cat. No.:	HY-P78220
Synonyms:	CD262; DR5; DR5TRICK2B; Fas-like protein; KILLER/DR5; TNFRSF10B; TRAIL R2; TRAIL receptor 2; TRAILR2; TRICK2; TRICK2A; TRICKB; ZTNFR9; KILLER
Species:	Human
Source:	HEK293
Accession:	O14763 (I56-E182)
Gene ID:	8795
Molecular Weight:	20-25 kDa

PROPERTIES

Biological Activity	Measured by its binding ability in a functional ELISA. When immobilized Anti-Trail R2 Antibody hFc Tag at 5 µg/mL (100 µl/Well), can bind Biotinylated Human TRAIL R2 His tag and the EC ₅₀ is 0.15 µg/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>The TRAIL R2/TNFRSF10B Protein functions as a receptor for the cytotoxic ligand TNFSF10/TRAIL. Upon ligand binding, the adapter molecule FADD recruits caspase-8 to the activated receptor, forming the death-inducing signaling complex (DISC), which triggers caspase-8 proteolytic activation and initiates the subsequent cascade of caspases, mediating apoptosis. Additionally, TRAIL R2/TNFRSF10B promotes the activation of NF-kappa-B and is essential for endoplasmic reticulum (ER) stress-induced apoptosis. In its monomeric state, it can interact with TRADD and RIPK1, and in the absence of stimulation, it interacts with BIRC2, DDX3X, and GSK3B. Stimulation of the receptor enhances interactions with BIRC2 and DDX3X, accompanied by their cleavage. Notably, TRAIL R2/TNFRSF10B can also interact with the HCMV protein UL141, preventing cell surface expression, where two TNFRSF10B monomers interact with a UL141 homodimer, and three TNFRSF10B molecules interact with TNFSF10 homotrimer. These intricate interactions underline the multifaceted role of TRAIL R2/TNFRSF10B in apoptotic and signaling pathways.</p>
-------------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA