

CD40 Protein, Cynomolgus (HEK 293, Fc)

Cat. No.:	HY-P7837
Synonyms:	rCynB-cell surface antigen CD40, Fc; Tumor Necrosis Factor Receptor Superfamily member 5; B-Cell Surface Antigen CD40; Bp50; CD40L Receptor; CDw40; CD40; TNFRSF5
Species:	Cynomolgus
Source:	HEK 293
Accession:	G7PG38 (E21-R193)
Gene ID:	102118696
Molecular Weight:	30-60 kDa

PROPERTIES

AA Sequence	<pre> E P P T A C R E K Q Y L I N S Q C C S L C Q P G Q K L V S D C T E F T E T E C L P C G E S E F L D T W N R E T R C H Q H K Y C D P N L G L R V Q Q K G T S E T D T I C T C E E G L H C T S E S C E S C V P H R S C L P G F G V K Q I A T G V S D T I C E P C P V G F F S N V S S A F E K C R P W T S C E T K D L V V Q Q A G T N K T D V V C G P Q D R Q R </pre>
Biological Activity	Data is not available.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 50 mM Tris-HCl, 100 mM Glycine, pH 7.5.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA