

## CD40 Protein, Cynomolgus (HEK293, Fc)

<b>Cat. No.:</b>	HY-P7837
<b>Synonyms:</b>	rCynB-cell surface antigen CD40, Fc; Tumor Necrosis Factor Receptor Superfamily member 5; B-Cell Surface Antigen CD40; Bp50; CD40L Receptor; CDw40; CD40; TNFRSF5
<b>Species:</b>	Cynomolgus
<b>Source:</b>	HEK293
<b>Accession:</b>	G7PG38 (E21-R193)
<b>Gene ID:</b>	102118696
<b>Molecular Weight:</b>	30-60 kDa

### PROPERTIES

<b>AA Sequence</b>	<pre> E P P T A C R E K Q   Y L I N S Q C C S L   C Q P G Q K L V S D   C T E F T E T E C L P C G E S E F L D T   W N R E T R C H Q H   K Y C D P N L G L R   V Q Q K G T S E T D T I C T C E E G L H   C T S E S C E S C V   P H R S C L P G F G   V K Q I A T G V S D T I C E P C P V G F   F S N V S S A F E K   C R P W T S C E T K   D L V V Q Q A G T N K T D V V C G P Q D   R Q R           </pre>
<b>Appearance</b>	Lyophilized powder.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 50 mM Tris-HCl, 100 mM Glycine, pH 7.5.
<b>Endotoxin Level</b>	<1 EU/µg, determined by LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
<b>Storage &amp; Stability</b>	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA