

B7-H4 Protein, Human (HEK293, His-Avi)

Cat. No.:	HY-P78390
Synonyms:	B7H4; B7-H4; B7S1; B7h.5; B7x; FLJ22418; VTCN1; PRO1291
Species:	Human
Source:	HEK293
Accession:	Q7Z7D3 (F29-A258)
Gene ID:	79679
Molecular Weight:	52-68 kDa

PROPERTIES

Biological Activity	Immobilized Human B7-H4, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-B7-H4 Antibody, hFc Tag with the EC ₅₀ of 9.4ng/ml determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	B7-H4 protein functions as a negative regulator of the T-cell-mediated immune response, exerting inhibitory effects on T-cell activation, proliferation, cytokine production, and the development of cytotoxicity. Particularly significant when expressed on the cell surface of tumor macrophages, B7-H4, in collaboration with regulatory T-cells (Treg), plays a crucial role in suppressing tumor-associated antigen-specific T-cell immunity. Additionally, B7-H4 is implicated in the promotion of epithelial cell transformation, indicating its multifaceted involvement in modulating immune responses and cellular processes associated with tumor development.
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA