

Siglec-10 Protein, Human (HEK293, Fc-Avi)

Cat. No.:	HY-P78516
Synonyms:	SLG2; SIGLEC10; MGC126774; PRO940
Species:	Human
Source:	HEK293
Accession:	Q96LC7 (M17-T546)
Gene ID:	89790
Molecular Weight:	95-110 kDa

PROPERTIES

Biological Activity	Immobilized Human Siglec-10 at 2µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Anti-Siglec-10 Antibody, hFc Tag with the EC ₅₀ of 14.4ng/ml determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

Siglec-10 protein, recognized as a putative adhesion molecule, functions in sialic-acid dependent cellular binding, displaying a preference for alpha-2,3- or alpha-2,6-linked sialic acid. The sialic acid recognition site of Siglec-10 may undergo masking due to cis interactions with sialic acids on the same cell surface. In immune responses, it appears to act as an inhibitory receptor, inducing ligand-induced tyrosine phosphorylation and recruiting cytoplasmic phosphatases via their SH2 domains, blocking signal transduction through dephosphorylation of signaling molecules. Siglec-10 is involved in the negative regulation of B-cell antigen receptor signaling, dependent on PTPN6/SHP-1. In association with CD24, it may participate in the selective suppression of the immune response to danger-associated molecular patterns (DAMPs) such as HMGB1, HSP70, and HSP90. Siglec-10, in collaboration with CD24, regulates the immune response of natural killer (NK) cells and plays a role in the control of autoimmunity. During the initiation of adaptive immune responses by CD8-alpha(+) dendritic cells, Siglec-10 inhibits cross-presentation by impairing the formation of MHC class I-peptide complexes, implicating the recruitment of PTPN6/SHP-1 and promoting phagosomal acidification. Siglec-10 interacts with various proteins, including PTPN6/SHP-1, NCF1, CD24, HMGB1, RIGI, CBL, and PTPN11.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA