

GMP IL-2 Protein, Human (solution)

Cat. No.:	HY-P78551
Synonyms:	Interleukin-2; IL-2; T-Cell Growth Factor; TCGF; Aldesleukin; IL2
Species:	Human
Source:	E. coli
Accession:	P60568 (A21-T153)
Gene ID:	3558
Molecular Weight:	Approximately 15.5 kDa

PROPERTIES

Biological Activity	Measured in a cell proliferation assay using CTLL2 and the ED ₅₀ for this effect is typically 1-8 ng/mL.
Appearance	Solution.
Formulation	Supplied as a 0.22 µm filtered solution of 20mM HAc-NaAc, 0.004% Tween-80, pH 4.0
Endotoxin Level	<0.005 EU/µg, determined by LAL method.
Reconstitution	N/A.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice.

DESCRIPTION

Background

The cytokine interleukin-2 (IL-2), produced primarily by activated CD4-positive helper T-cells and, to a lesser extent, by activated CD8-positive T-cells and natural killer (NK) cells, plays pivotal roles in the immune response and tolerance. IL-2 binds to a receptor complex composed of either the high-affinity trimeric IL-2R (IL2RA/CD25, IL2RB/CD122, and IL2RG/CD132) or the low-affinity dimeric IL-2R (IL2RB and IL2RG). This interaction induces oligomerization and conformational changes in the IL-2R subunits, initiating downstream signaling with the phosphorylation of JAK1 and JAK3. Subsequently, JAK1 and JAK3 phosphorylate the receptor, creating a docking site for the phosphorylation of various substrates, including STAT5. This process activates multiple pathways, including STAT, phosphoinositide-3-kinase/PI3K, and mitogen-activated protein kinase/MAPK pathways. IL-2 functions as a T-cell growth factor, enhances NK-cell cytolytic activity, and promotes robust proliferation of activated B-cells, leading to increased immunoglobulin production. Furthermore, IL-2 plays a crucial role in regulating the adaptive immune system by controlling the survival and proliferation of regulatory T-cells, essential for maintaining immune tolerance. Additionally, IL-2 participates in the differentiation and homeostasis of various effector T-cell subsets, including Th1, Th2, Th17, as well as memory CD8-positive T-cells.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA