

gp130/IL6ST Protein, Mouse (595a.a, HEK293, His)

Cat. No.:	HY-P78781
Synonyms:	IL6ST; gp130; CD130; IL-6RB; IL-6R-beta; CDw130
Species:	Mouse
Source:	HEK293
Accession:	NP_034690/Q00560 (Q23-E617)
Gene ID:	16195
Molecular Weight:	80-100 kDa

PROPERTIES

AA Sequence

Q L L E P C G Y I Y	P E F P V V Q R G S	N F T A I C V L K E	A C L Q H Y Y V N A
S Y I V W K T N H A	A V P R E Q V T V I	N R T T S S V T F T	D V V L P S V Q L T
C N I L S F G Q I E	Q N V Y G V T M L S	G F P P D K P T N L	T C I V N E G K N M
L C Q W D P G R E T	Y L E T N Y T L K S	E W A T E K F P D C	Q S K H G T S C M V
S Y M P T Y Y V N I	E V W V E A E N A L	G K V S S E S I N F	D P V D K V K P T P
P Y N L S V T N S E	E L S S I L K L S W	V S S G L G G L L D	L K S D I Q Y R T K
D A S T W I Q V P L	E D T M S P R T S F	T V Q D L K P F T E	Y V F R I R S I K D
S G K G Y W S D W S	E E A S G T T Y E D	R P S R P P S F W Y	K T N P S H G Q E Y
R S V R L I W K A L	P L S E A N G K I L	D Y E V I L T Q S K	S V S Q T Y T V T G
T E L T V N L T N D	R Y V A S L A A R N	K V G K S A A A V L	T I P S P H V T A A
Y S V V N L K A F P	K D N L L W V E W T	P P P K P V S K Y I	L E W C V L S E N A
P C V E D W Q Q E D	A T V N R T H L R G	R L L E S K C Y Q I	T V T P V F A T G P
G G S E S L K A Y L	K Q A A P A R G P T	V R T K K V G K N E	A V L A W D Q I P V
D D Q N G F I R N Y	S I S Y R T S V G K	E M V V H V D S S H	T E Y T L S S L S S
D T L Y M V R M A A	Y T D E G G K D G P	E F T F T T P K F A	Q G E I E

Biological Activity

1. Measured by its ability to inhibit IL-6-dependent proliferation of TF-1 cells. The ED₅₀ this effect is 0.1185 ng/mL in the presence of 4 µg/mL Recombinant Human IL-6 R alpha , corresponding to a specific activity is 8.44×10⁶ units/mg.
 2. Immobilized Human gp130, His Tag at 5 µg/mL (100 µL/well) can bind Biotinylated Human IL-6 R alpha that produces 50% of the optimal binding response is found to be approximately 6.8 ng/mL.

Appearance

Lyophilized powder

Formulation

Lyophilized a 0.22 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.

Endotoxin Level

<1 EU/µg, determined by LAL method.

Reconstitution

It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH₂O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).

Storage & Stability

Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is

recommended to freeze aliquots at -20°C or -80°C for extended storage.

Shipping

Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

Glycoprotein 130 mRNA is ubiquitously expressed in the human body, exhibiting highest levels in saphenous vein, pericardium, ovary, omental adipose tissue, peritoneum, spleen lymphnodes, and trigeminal ganglia. And gp130 expression is significantly increased in adult organs compared to fetal organs. Moreover, gp130 is highly expressed in multiple types of cancers compared to normal tissues^[3].

The amino acid sequence of human gp130 protein has low homology between mouse and rat gp130 protein.

gp130 signaling is initiated when autocrine or paracrine cyto-kines induce the dimerization of gp130 and an α -receptor subunit to form the receptor complex, leading to the activation of downstream signaling cascades, such as the JAK/STAT, the PI3K/Akt, and the Ras/Raf/Mek/Erk1/2/MAPK pathways. The activation of JAK/Stat3 pathway induces its target gene expression, resulting in cell growth and proliferation (cyclin D, CDC25 A, c-Myc, survivin), survival (survivin, Bcl-2, Bcl-X_L), angiogenesis (VEGF), and cell migration (MMP-2, MMP-7, and MMP-9)^[3]. gp130 is a IL-6 antagonist shows anti-tumor, anti-inflammation and antinociceptive activity^[4]. gp130 decreases the STAT3 phosphorylation induced by IL-6, and decreases IL-6-induced increase of Ki67^[5].

REFERENCES

- [1]. Cao Y, et al. Glycoprotein 130 is associated with adverse postoperative clinical outcomes of patients with late-stage non-metastatic gastric cancer. *Sci Rep.* 2016 Dec 5;6:38364.
- [2]. Nogueira-Silva C, et al. The role of glycoprotein 130 family of cytokines in fetal rat lung development. *PLoS One.* 2013 Jun 24;8(6):e67607.
- [3]. Xu S, et al. gp130: a promising drug target for cancer therapy. *Expert Opin Ther Targets.* 2013 Nov;17(11):1303-28.
- [4]. Boettger MK, et al. Differential effects of locally and systemically administered soluble glycoprotein 130 on pain and inflammation in experimental arthritis. *Arthritis Res Ther.* 2010;12(4):R140.
- [5]. Hong J, et al. Recombinant soluble gp130 protein reduces DEN-induced primary hepatocellular carcinoma in mice. *Sci Rep.* 2016 Apr 15;6:24397.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA