

GM-CSF Protein, Human (Tag Free)

Cat. No.:	HY-P78868
Synonyms:	GM-CSF; CSF2; MGC131935
Species:	Human
Source:	HEK293
Accession:	P04141 (A18-E144)
Gene ID:	1437
Molecular Weight:	Approximately 19-29 kDa due to the glycosylation

PROPERTIES

AA Sequence	<p>A P A R S P S P S T Q P W E H V N A I Q E A R R L L N L S R D T A A E M N E T V</p> <p>E V I S E M F D L Q E P T C L Q T R L E L Y K Q G L R G S L T K L K G P L T M M</p> <p>A S H Y K Q H C P P T P E T S C A T Q I I T F E S F K E N L K D F L L V I P F D</p> <p>C W E P V Q E</p>
Biological Activity	Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED ₅₀ for this effect is ≤ 1.731 ng/mL, corresponding to a specific activity is ≥ 5.777×10 ⁵ units/mg.
Appearance	Lyophilized powder
Formulation	Lyophilized from a 0.22 μm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4 or PBS, pH 7.4.
Endotoxin Level	<1 EU/μg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	GMP GM-CSF Protein functions as a pivotal cytokine, fostering the growth and differentiation of hematopoietic precursor cells spanning diverse lineages, such as granulocytes, macrophages, eosinophils, and erythrocytes. In its monomeric form, GMP GM-CSF serves as a signaling molecule, orchestrating a complex receptor assembly. This receptor complex takes the shape of a dodecamer, consisting of two head-to-head hexamers, each composed of two alpha, two beta, and two ligand subunits. This structural intricacy underscores the specificity and regulatory role of GMP GM-CSF in directing cellular
------------	---

responses within the hematopoietic system.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA