

## Product Data Sheet

### ASC Antibody

| Cat. No.:         | HY-P80548   |  |
|-------------------|---|--|
| Synonyms:         | ASC Antibody is a non-conjugated and Rabbit origined polyclonal antibody about 22 kDa,<br>targeting to ASC. It can be used for WB,IHC-F,IHC-P,ICC/IF,ELISA assays with tag free, in the |  |
|                   | background of Human, Mouse, Rat.  |  |
| Host:             | Rabbit  |  |
| Reactivity:       | Human,Mouse,Rat   |  |
| Conjugation:      | Non-conjugated  |  |
| SwissProt ID:     | Q9ULZ3  |  |
| Research Field:   | Cell Biology  |  |
| Molecular Weight: | Predicted band size: 22 kDa   |  |

| PROPERTIES                      |  |                |  |
|---------------------------------|--|----------------|--|
| Formulation                     | Supplied in 1*PBS (pH 7.3), 50% glycerol and 0.5% BSA. Preservative: 0.02% sodium azide. |                |  |
| Purity                          | affinity purified  |                |  |
| Storage & Stability             | Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.                         |                |  |
| Appearance                      | Liquid   |                |  |
| Application &<br>Dilution Ratio | Application  | Dilution Ratio |  |
|                                 | WB   | 1:500-1:1,000  |  |
|                                 | ІНС  | 1:50-1:100     |  |
|                                 | IF   | 1:50-1:200     |  |
|                                 | ELISA  | 1:10,000       |  |
| Shipping                        | Shipping with blue ice.  |                |  |

#### DESCRIPTION

# BackgroundASC: This gene encodes an adaptor protein that is composed of two protein-protein interaction domains: a N-terminal<br/>PYRIN-PAAD-DAPIN domain (PYD) and a C-terminal caspase-recruitment domain (CARD). The PYD and CARD domains are<br/>members of the six-helix bundle death domain-fold superfamily that mediates assembly of large signaling complexes in the<br/>inflammatory and apoptotic signaling pathways via the activation of caspase. In normal cells, this protein is localized to the<br/>cytoplasm; however, in cells undergoing apoptosis, it forms ball-like aggregates near the nuclear periphery. Two transcript<br/>variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

#### Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA