

## Glutathione Peroxidase 4 Antibody

Cat. No.:	HY-P80692
Synonyms:	Glutathione Peroxidase 4 Antibody is a non-conjugated and Rabbit originated polyclonal antibody about 22 kDa, targeting to Glutathione Peroxidase 4. It can be used for WB,IHC-F,IHC-P,ICC/IF assays with tag free, in the background of Human, Mouse, Rat.
Host:	Rabbit
Reactivity:	Human,Mouse,Rat
Conjugation:	Non-conjugated
SwissProt ID:	P36969
Research Field:	Signal Transduction
Molecular Weight:	Predicted band size: 22 kDa

### PROPERTIES

Formulation	Supplied in phosphate buffered saline (pH 7.4), 150 mM NaCl and 50% glycerol. Preservative: 0.02% sodium azide	
Purity	affinity purified	
Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.	
Appearance	Liquid	
Application & Dilution Ratio	Application	Dilution Ratio
	WB	1:500-1:1,000
	IHC	1:50-1:100
	IF	1:50-1:200
Shipping	Shipping with blue ice.	

### DESCRIPTION

#### Background

Glutathione Peroxidase 4: The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyze the reduction of hydrogen peroxide, organic hydroperoxides and lipid hydroperoxides, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity. This isozyme has a high preference for lipid hydroperoxides and protects cells against membrane lipid peroxidation and cell death. It is also required for normal sperm development; thus, it has been identified as a 'moonlighting' protein because of its ability to serve dual functions as a peroxidase, as well as a structural protein in mature spermatozoa. Mutations in this gene are associated with Sedaghatian type of spondylometaphyseal dysplasia (SMDS). This isozyme is also a selenoprotein, containing the rare amino acid selenocysteine (Sec) at its active site. Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Transcript variants resulting from alternative splicing or use of alternate promoters have

---

been described to encode isoforms with different subcellular localization. [provided by RefSeq, Dec 2018]

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA