**Proteins** 

# **Screening Libraries**

# **Product** Data Sheet



## **IL-1** beta Antibody

HY-P80720 Cat. No.:

Synonyms: IL-1 beta Antibody is a non-conjugated and Rabbit origined polyclonal antibody about 31

kDa, targeting to IL-1 beta. It can be used for WB,IHC-P,ICC/IF assays with tag free, in the

background of Human, Mouse, Rat.

Host: Rabbit

Reactivity: Human, Mouse, Rat Conjugation: Non-conjugated

SwissProt ID: P01584

Research Field: **Immunology** 

Predicted band size: 31 kDa Molecular Weight:

### **PROPERTIES**

FC	rı	nu	la	tio	n

Supplied in 1\*PBS (pH 7.3), 50% glycerol and 0.5% BSA. Preservative: 0.02% sodium azide.

Purity

affinity purified

Storage & Stability

Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.

**Appearance** 

Liquid

Application &				
<b>Dilution Ratio</b>				

Application	Dilution Ratio
WB	1:500-1:1,000
IHC	1:50-1:100
IF	1:50-1:200

**Shipping** 

Shipping with blue ice.

### **DESCRIPTION**

### Background

IL-1 beta: The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. Similarly, IL-1B has been implicated in human osteoarthritis pathogenesis. Patients with severe Coronavirus Disease 2019 (COVID-19) present elevated levels of pro-inflammatory cytokines such as IL-1B in bronchial alveolar lavage fluid samples. The lung damage induced by the Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL-1B. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, Jul 2020]

Page 1 of 2

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$ 

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com