

Lactate Dehydrogenase C Antibody

Cat. No.:	HY-P80739
Synonyms:	Lactate Dehydrogenase C Antibody is a non-conjugated and Rabbit originated monoclonal antibody about 36 kDa, targeting to Lactate Dehydrogenase C. It can be used for WB assays with tag free, in the background of Human.
Host:	Rabbit
Reactivity:	Human
Conjugation:	Non-conjugated
SwissProt ID:	P07864
Research Field:	Signal Transduction
Molecular Weight:	Predicted band size: 36 kDa

PROPERTIES

Formulation	Supplied in 50 mM Tris-Glycine (pH 7.4), 0.15 M NaCl, 40% Glycerol and 0.05% BSA. Preservative: 0.01% Sodium azide					
Purity	affinity purified					
Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.					
Appearance	Liquid					
Application & Dilution Ratio	<table border="1"> <thead> <tr> <th>Application</th> <th>Dilution Ratio</th> </tr> </thead> <tbody> <tr> <td>WB</td> <td>1:500-1:1,000</td> </tr> </tbody> </table>	Application	Dilution Ratio	WB	1:500-1:1,000	
Application	Dilution Ratio					
WB	1:500-1:1,000					
Shipping	Shipping with blue ice.					

DESCRIPTION

Background	Lactate Dehydrogenase C: Lactate dehydrogenase C catalyzes the conversion of L-lactate and NAD to pyruvate and NADH in the final step of anaerobic glycolysis. LDHC is testis-specific and belongs to the lactate dehydrogenase family. Two transcript variants have been detected which differ in the 5' untranslated region. [provided by RefSeq, Jul 2008]
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA