

Product Data Sheet

LRP1 Antibody

Cat. No.:	HY-P80746	
Synonyms:	LRP1 Antibody is a non-conjugated and Rabbit origined monoclonal antibody about 505 kDa,	
	targeting to LRP1. It can be used for WB,IHC-F,IHC-P,ICC/IF,IP assays with tag free, in the	
	background of Human, Mouse, Rat.	
Host:	Rabbit	
Reactivity:	Human,Mouse,Rat	
Conjugation:	Non-conjugated	
SwissProt ID:	Q07954	
Research Field:	Neuroscience	
Molecular Weight:	Predicted band size: 85 kDa	

Formulation Supplied in 50 mM Tris-Glycine (pH 7.4), 0.15 M NaCl, 40% Glycerol and 0.05% BSA. Preservative: 0.01% Sodium azide Purity affinity purified Storage & Stability Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles. Appearance Liquid Application & Dilution Ratio Dilution Ratio IDIUtion Ratio 1:500-1:1,000 IHC 1:50-1:200 IF 1:50-1:200 IP 1:20	PROPERTIES			
Storage & Stability Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles. Appearance Liquid Application & Dilution Ratio Dilution Ratio IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII		Supplied in 50 mM Tris-Glycine (pH 7.4), 0.15 M NaCl, 40% Glycerol and 0.05% BSA. Preservative: 0.01% Sodium azide		
Appearance Liquid Application & Dilution Ratio Application MB 1:500-1:1,000 IHC 1:50-1:100 IF 1:50-1:200 IP 1:20	Purity	affinity purified		
Application & Dilution RatioApplicationDilution RatioWB1:500-1:1,000IHC1:50-1:100IF1:50-1:200IP1:20	Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.		
Dilution Ratio WB 1:500-1:1,000 IHC 1:50-1:100 IF 1:50-1:200 IP 1:20	Appearance	Liquid		
Instant Instant Instant INC 1:50-1:100 IF 1:50-1:200 IP 1:20	Application &	Application	Dilution Ratio	
IF 1:50-1:200 IP 1:20	Dilution Ratio	WB	1:500-1:1,000	
IP 1:20		ІНС	1:50-1:100	
		IF	1:50-1:200	
Shipping Shipping with blue ice.		IP	1:20	
	Shipping	Shipping with blue ice.		

DESCRIPTION

Background

LRP1: This gene encodes a member of the low-density lipoprotein receptor family of proteins. The encoded preproprotein is proteolytically processed by furin to generate 515 kDa and 85 kDa subunits that form the mature receptor (PMID: 8546712). This receptor is involved in several cellular processes, including intracellular signaling, lipid homeostasis, and clearance of apoptotic cells. In addition, the encoded protein is necessary for the alpha 2-macroglobulin-mediated clearance of secreted amyloid precursor protein and beta-amyloid, the main component of amyloid plaques found in Alzheimer patients. Expression of this gene decreases with age and has been found to be lower than controls in brain tissue from Alzheimer's disease patients. [provided by RefSeq, Oct 2015]

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA