

## MSR1 Antibody

Cat. No.:	HY-P80758
Synonyms:	MSR1 Antibody is a non-conjugated and Rabbit originated polyclonal antibody about 50 kDa, targeting to MSR1. It can be used for WB assays with tag free, in the background of Human, Mouse, Rat.
Host:	Rabbit
Reactivity:	Human, Mouse, Rat
Conjugation:	Non-conjugated
SwissProt ID:	P21757
Research Field:	Immunology
Molecular Weight:	Predicted band size: 50 kDa

### PROPERTIES

Formulation	Supplied in phosphate buffered saline (pH 7.4), 150 mM NaCl and 50% glycerol. Preservative: 0.02% sodium azide					
Purity	affinity purified					
Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.					
Appearance	Liquid					
Application & Dilution Ratio	<table> <thead> <tr> <th>Application</th> <th>Dilution Ratio</th> </tr> </thead> <tbody> <tr> <td>WB</td> <td>1:500-1:1,000</td> </tr> </tbody> </table>	Application	Dilution Ratio	WB	1:500-1:1,000	
Application	Dilution Ratio					
WB	1:500-1:1,000					
Shipping	Shipping with blue ice.					

### DESCRIPTION

Background	<p>MSR1: This gene encodes the class A macrophage scavenger receptors, which include three different types (1, 2, 3) generated by alternative splicing of this gene. These receptors or isoforms are macrophage-specific trimeric integral membrane glycoproteins and have been implicated in many macrophage-associated physiological and pathological processes including atherosclerosis, Alzheimer's disease, and host defense. The isoforms type 1 and type 2 are functional receptors and are able to mediate the endocytosis of modified low density lipoproteins (LDLs). The isoform type 3 does not internalize modified LDL (acetyl-LDL) despite having the domain shown to mediate this function in the types 1 and 2 isoforms. It has an altered intracellular processing and is trapped within the endoplasmic reticulum, making it unable to perform endocytosis. The isoform type 3 can inhibit the function of isoforms type 1 and type 2 when co-expressed, indicating a dominant negative effect and suggesting a mechanism for regulation of scavenger receptor activity in macrophages. [provided by RefSeq, Jul 2008]</p>
------------	--

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA