

NLRP3 Antibody

Cat. No.:	HY-P80766
Synonyms:	NLRP3 Antibody is a non-conjugated and Rabbit originated polyclonal antibody about 118 kDa, targeting to NLRP3. It can be used for WB,IHC-P,ICC/IF,IP,FC assays with tag free, in the background of Mouse, Rat.
Host:	Rabbit
Reactivity:	Human,Mouse,Rat
Conjugation:	Non-conjugated
SwissProt ID:	Q96P20
Research Field:	Immunology
Molecular Weight:	Predicted band size: 118 kDa

PROPERTIES

Formulation	Supplied in phosphate buffered saline (pH 7.4), 150 mM NaCl and 50% glycerol. Preservative: 0.02% sodium azide	
Purity	affinity purified	
Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.	
Appearance	Liquid	
Application & Dilution Ratio	Application	Dilution Ratio
	WB	1:500-1:1,000
	IHC	1:50-1:100
	IF	1:50-1:200
	IP	1:20
	FC	1:50-1:100
Shipping	Shipping with blue ice.	

DESCRIPTION

Background	<p>NLRP3: This gene encodes a pyrin-like protein containing a pyrin domain, a nucleotide-binding site (NBS) domain, and a leucine-rich repeat (LRR) motif. This protein interacts with the apoptosis-associated speck-like protein PYCARD/ASC, which contains a caspase recruitment domain, and is a member of the NLRP3 inflammasome complex. This complex functions as an upstream activator of NF-kappaB signaling, and it plays a role in the regulation of inflammation, the immune response, and apoptosis. The SARS-CoV 3a protein, a transmembrane pore-forming viroporin, has been shown to activate the NLRP3 inflammasome via the formation of ion channels in macrophages. Mutations in this gene are associated with familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome, neonatal-onset multisystem inflammatory disease (NOMID), keratoendotheliitis fugax hereditarian, and</p>
------------	--

deafness, autosomal dominant 34, with or without inflammation. Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Alternative 5' UTR structures are suggested by available data; however, insufficient evidence is available to determine if all of the represented 5' UTR splice patterns are biologically valid. [provided by RefSeq, Aug 2020]

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA