

Product Data Sheet

Nrf1 Antibody

HY-P80769
Nrf1 Antibody is a non-conjugated and Rabbit origined monoclonal antibody about 54 kDa, targeting to Nrf1. It can be used for WB,IHC-F,IHC-P,ICC/IF,IP assays with tag free, in the background of Human, Mouse, Rat.
Rabbit
Human,Mouse,Rat
Non-conjugated
Q16656
Epigenetics and Nuclear Signaling
Predicted band size: 54 kDa

Formulation Supplied in 50 mM Tris-Glycine (pH 7.4), 0.15 M NaCl, 40% Glycerol and 0.05% BSA. Preservative: 0.01% Sodium azide Purity affinity purified Storage & Stability Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles. Appearance Liquid Application & Dilution Ratio Dilution Ratio III HC 1:500-1:1,000 IF 1:50-1:200 IP 1:20	PROPERTIES		
Storage & Stability Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles. Appearance Liquid Application & Dilution Ratio Dilution Ratio IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII		Supplied in 50 mM Tris-Glycine (pH 7.4), 0.15 M NaCl, 40% Glycerol and 0.05% BSA. Preservative: 0.01% Sodium azide	
Appearance Liquid Application & Dilution Ratio Application MB 1:500-1:1,000 IHC 1:50-1:100 IF 1:50-1:200 IP 1:20	Purity	affinity purified	
Application & Dilution RatioApplicationDilution RatioWB1:500-1:1,000IHC1:50-1:100IF1:50-1:200IP1:20	Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.	
Dilution Ratio WB 1:500-1:1,000 IHC 1:50-1:100 IF 1:50-1:200 IP 1:20	Appearance	Liquid	
Instant Instant Instant INC 1:50-1:100 IF 1:50-1:200 IP 1:20	Application &	Application	Dilution Ratio
IF 1:50-1:200 IP 1:20	Dilution Ratio	WB	1:500-1:1,000
IP 1:20		ІНС	1:50-1:100
		IF	1:50-1:200
Shipping Shipping with blue ice.		IP	1:20
	Shipping	Shipping with blue ice.	

DESCRIPTION

Background

Nrf1: This gene encodes a protein that homodimerizes and functions as a transcription factor which activates the expression of some key metabolic genes regulating cellular growth and nuclear genes required for respiration, heme biosynthesis, and mitochondrial DNA transcription and replication. The protein has also been associated with the regulation of neurite outgrowth. Alternative splicing results in multiple transcript variants. Confusion has occurred in bibliographic databases due to the shared symbol of NRF1 for this gene and for "nuclear factor (erythroid-derived 2)-like 1" which has an official symbol of NFE2L1. [provided by RefSeq, May 2014]

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA