

p38 Antibody

Cat. No.:	HY-P80777
Synonyms:	p38 Antibody is a non-conjugated and Rabbit originated monoclonal antibody about 41 kDa, targeting to p38. It can be used for WB,IP assays with tag free, in the background of Human.
Host:	Rabbit
Reactivity:	Human
Conjugation:	Non-conjugated
SwissProt ID:	Q16539
Research Field:	Signal Transduction
Molecular Weight:	Predicted band size: 41 kDa

PROPERTIES

Formulation	Supplied in 50 mM Tris-Glycine (pH 7.4), 0.15 M NaCl, 40% Glycerol and 0.05% BSA. Preservative: 0.01% Sodium azide	
Purity	affinity purified	
Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.	
Appearance	Liquid	
Application & Dilution Ratio	Application	Dilution Ratio
	WB	1:500-1:1,000
	IP	1:20
Shipping	Shipping with blue ice.	

DESCRIPTION

Background	<p>p38: The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]</p>
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA