

Phospho-c-Myc (Ser62) Antibody

Cat. No.:	HY-P80804
Synonyms:	Phospho-c-Myc (Ser62) Antibody is a non-conjugated and Rabbit originated monoclonal antibody about 49 kDa, targeting to Phospho-c-Myc (Ser62). It can be used for WB, ICC/IF, IP assays with tag free, in the background of Mouse, Rat.
Host:	Rabbit
Reactivity:	Mouse, Rat
Conjugation:	Non-conjugated
SwissProt ID:	P01106
Research Field:	Epigenetics and Nuclear Signaling
Molecular Weight:	Predicted band size: 49 kDa

PROPERTIES

Formulation	Supplied in 50 mM Tris-Glycine (pH 7.4), 0.15 M NaCl, 40% Glycerol and 0.05% BSA. Preservative: 0.01% Sodium azide	
Purity	affinity purified	
Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.	
Appearance	Liquid	
Application & Dilution Ratio	Application	Dilution Ratio
	WB	1:500-1:1,000
	IF	1:50-1:200
	IP	1:20
Shipping	Shipping with blue ice.	

DESCRIPTION

Background	<p>c-Myc: This gene is a proto-oncogene and encodes a nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. The encoded protein forms a heterodimer with the related transcription factor MAX. This complex binds to the E box DNA consensus sequence and regulates the transcription of specific target genes. Amplification of this gene is frequently observed in numerous human cancers. Translocations involving this gene are associated with Burkitt lymphoma and multiple myeloma in human patients. There is evidence to show that translation initiates both from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site, resulting in the production of two isoforms with distinct N-termini. [provided by RefSeq, Aug 2017]</p>
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA