

Phospho-eIF2A (Ser51) Antibody

Cat. No.:	HY-P80811
Synonyms:	Phospho-eIF2A (Ser51) Antibody is a non-conjugated and Rabbit originated monoclonal antibody about 36 kDa, targeting to Phospho-eIF2A (Ser51). It can be used for WB assays with tag free, in the background of Human, Mouse, Rat.
Host:	Rabbit
Reactivity:	Human, Mouse, Rat
Conjugation:	Non-conjugated
SwissProt ID:	P05198
Research Field:	Epigenetics and Nuclear Signaling
Molecular Weight:	Predicted band size: 36 kDa

PROPERTIES

Formulation	Supplied in 50 mM Tris-Glycine (pH 7.4), 0.15 M NaCl, 40% Glycerol and 0.05% BSA. Preservative: 0.01% Sodium azide					
Purity	affinity purified					
Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.					
Appearance	Liquid					
Application & Dilution Ratio	<table border="1"> <thead> <tr> <th>Application</th> <th>Dilution Ratio</th> </tr> </thead> <tbody> <tr> <td>WB</td> <td>1:500-1:1,000</td> </tr> </tbody> </table>	Application	Dilution Ratio	WB	1:500-1:1,000	
Application	Dilution Ratio					
WB	1:500-1:1,000					
Shipping	Shipping with blue ice.					

DESCRIPTION

Background	eIF2A: The translation initiation factor EIF2 catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. Binding occurs as a ternary complex of methionyl-tRNA, EIF2, and GTP. EIF2 is composed of 3 nonidentical subunits, the 36-kD EIF2-alpha subunit (EIF2S1), the 38-kD EIF2-beta subunit (EIF2S2; MIM 603908), and the 52-kD EIF2-gamma subunit (EIF2S3; MIM 300161). The rate of formation of the ternary complex is modulated by the phosphorylation state of EIF2-alpha (Ernst et al., 1987 [PubMed 2948954]).[supplied by OMIM, Feb 2010]
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA