

Thioredoxin Reductase 2 Antibody (YA661)

Cat. No.:	HY-P80913
Synonyms:	Thioredoxin Reductase 2 Antibody (YA661) is a non-conjugated and Mouse originated monoclonal antibody about 57 kDa, targeting to Thioredoxin Reductase 2 (3F2). It can be used for WB assays with tag free, in the background of Human.
Host:	Mouse
Reactivity:	Human
Conjugation:	Non-conjugated
SwissProt ID:	Q9NNW7
Research Field:	Cell Biology
Molecular Weight:	Predicted band size: 57 kDa

PROPERTIES

Formulation	Supplied in 1*PBS (pH 7.3), 50% glycerol and 0.5% BSA. Preservative: 0.02% sodium azide.					
Purity	affinity purified					
Storage & Stability	Stored at -20°C for 1 year. Avoid repeated freeze / thaw cycles.					
Appearance	Liquid					
Application & Dilution Ratio	<table> <thead> <tr> <th>Application</th> <th>Dilution Ratio</th> </tr> </thead> <tbody> <tr> <td>WB</td> <td>1:500-1:1,000</td> </tr> </tbody> </table>	Application	Dilution Ratio	WB	1:500-1:1,000	
Application	Dilution Ratio					
WB	1:500-1:1,000					
Shipping	Shipping with blue ice.					

DESCRIPTION

Background	<p>Thioredoxin Reductase 2 (3F2): The protein encoded by this gene belongs to the pyridine nucleotide-disulfide oxidoreductase family, and is a member of the thioredoxin (Trx) system. Three thioredoxin reductase (TrxR) isozymes are found in mammals. TrxRs are selenocysteine-containing flavoenzymes, which reduce thioredoxins, as well as other substrates, and play a key role in redox homeostasis. This gene encodes a mitochondrial form important for scavenging reactive oxygen species in mitochondria. It functions as a homodimer containing FAD, and selenocysteine (Sec) at the active site. Sec is encoded by UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, the Sec insertion sequence (SECIS) element, which is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. Alternatively spliced transcript variants encoding different isoforms, including a few localized in the cytosol and some lacking the C-terminal Sec residue, have been found for this gene. [provided by RefSeq, Jun 2017]</p>
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA