

## 3-Butynoic acid

Cat. No.: HY-W004784 CAS No.: 2345-51-9 Molecular Formula:  $C_4H_4O_2$  Molecular Weight: 84.07


Target: Endogenous Metabolite

Pathway: Metabolic Enzyme/Protease

Storage: Powder -20°C 3 years

4°C 2 years In solvent -80°C 6 months

-20°C 1 month



**Product** Data Sheet

## **SOLVENT & SOLUBILITY**

In Vitro H<sub>2</sub>O:

 $\rm H_2O$  : 100 mg/mL (1189.48 mM; Need ultrasonic)

DMSO : ≥ 100 mg/mL (1189.48 mM)

\* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	11.8948 mL	59.4743 mL	118.9485 mL
	5 mM	2.3790 mL	11.8948 mL	23.7897 mL
	10 mM	1.1895 mL	5.9474 mL	11.8948 mL

Please refer to the solubility information to select the appropriate solvent.

## **BIOLOGICAL ACTIVITY**

Description	3-Butynoic acid is an inhibitor of acyl CoA dehydrogenase <sup>[1]</sup> . 3-Butynoic acid is a click chemistry reagent, it contains an Alkyne group and can undergo copper-catalyzed azide-alkyne cycloaddition (CuAAc) with molecules containing Azide groups.
IC <sub>50</sub> & Target	Human Endogenous Metabolite

## **REFERENCES**

[1]. Akedo, M., et al. Direct Demonstration of Lactate-Acrylate Interconversion in Clostridium Propionicum. Nat Biotechnol 1, 791–794 (1983).

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$ 

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com