Boc-Ser(Me)-OH

MedChemExpress

| Cat. No.: | HY-W00802 | 2 | |
|--------------------|------------------------------------------------|----------|----------|
| CAS No.: | 51293-47-1 | | |
| Molecular Formula: | C ₉ H ₁₇ NO ₅ | | |
| Molecular Weight: | 219.23 | | |
| Target: | Amino Acid | Derivati | /es |
| Pathway: | Others | | |
| Storage: | Powder | -20°C | 3 years |
| | | 4°C | 2 years |
| | In solvent | -80°C | 6 months |
| | | -20°C | 1 month |
| | | | |

®

SOLVENT & SOLUBILITY

| | | Solvent Mass Concentration | 1 mg | 5 mg | 10 mg | | |
|---------|------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|--------------------|------------|------------|--|--|
| | Preparing Stock Solutions | 1 mM | 4.5614 mL | 22.8071 mL | 45.6142 mL | | |
| | | 5 mM | 0.9123 mL | 4.5614 mL | 9.1228 mL | | |
| | | 10 mM | 0.4561 mL | 2.2807 mL | 4.5614 mL | | |
| | Please refer to the so | lubility information to select the app | propriate solvent. | | | | |
| In Vivo | | 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (11.40 mM); Clear solution | | | | | |
| | | 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (11.40 mM); Clear solution | | | | | |
| | | vent one by one: 10% DMSO >> 90% corn oil 2.5 mg/mL (11.40 mM); Clear solution | | | | | |

| BIOLOGICAL ACTIVITY | | | | |
|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|
| Description | Boc-Ser(Me)-OH is a serine derivative ^[1] . | | | |
| In Vitro | Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. | | | |

Product Data Sheet

0

Ο

) NH

νOH

.0

REFERENCES

[1]. Luckose F, et al. Effects of amino acid derivatives on physical, mental, and physiological activities. Crit Rev Food Sci Nutr. 2015;55(13):1793-1144.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA