


Sinapinic acid

Cat. No.:	HY-W009732		
CAS No.:	530-59-6		
Molecular Formula:	C ₁₁ H ₁₂ O ₅		
Molecular Weight:	224.21		
Target:	HDAC; Angiotensin-converting Enzyme (ACE); Reactive Oxygen Species; Apoptosis		
Pathway:	Cell Cycle/DNA Damage; Epigenetics; Metabolic Enzyme/Protease; Immunology/Inflammation; NF-κB; Apoptosis		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (446.01 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM	4.4601 mL	22.3005 mL	44.6010 mL
		5 mM	0.8920 mL	4.4601 mL	8.9202 mL
10 mM		0.4460 mL	2.2301 mL	4.4601 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (11.15 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (11.15 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (11.15 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	<p>Sinapinic acid (Sinapic acid) is a phenolic compound isolated from <i>Hydnophytum formicarum</i> Jack. Rhizome, acts as an inhibitor of HDAC, with an IC₅₀ of 2.27 mM^[1], and also inhibits ACE-I activity^[2]. Sinapinic acid possess potent anti-tumor activity, induces apoptosis of tumor cells^[1]. Sinapinic acid shows antioxidant and antidiabetic activities^[2]. Sinapinic acid reduces total cholesterol, triglyceride, and HOMA-IR index, and also normalizes some serum</p>
-------------	---

	parameters of antioxidative abilities and oxidative damage in ovariectomized rats ^[3] .	
IC₅₀ & Target	HDAC 2.27 mM (IC ₅₀)	ACE-I
In Vitro	Sinapinic acid acts as an inhibitor of HDAC, with an IC ₅₀ of 2.27 mM ^[1] . Sinapinic acid also inhibits ACE-I activity ^[2] . Sinapinic acid inhibits HDAC activity in HeLa cells, suppresses the growth of HeLa and HT29 cells with IC ₅₀ s of 0.91 ± 0.02 mM and 1.6 ± 0.02 mM at 72 h, respectively, induces apoptosis of these cancer cells ^[1] .	
In Vivo	Sinapinic acid (5 or 25 mg/kg, p.o. daily for 4 weeks) increases the serum estradiol concentration; decreases insulin resistance and the triglyceride and total cholesterol concentrations; and favorably affects the parameters of antioxidant abilities (reduces glutathione, superoxide dismutase) and oxidative damage in rats ^[3] .	

REFERENCES

- [1]. Senawong T, et al. Histone deacetylase (HDAC) inhibitory and antiproliferative activities of phenolic-rich extracts derived from the rhizome of *Hydnophytum formicarum* Jack.: sinapinic acid acts as HDAC inhibitor. *BMC Complement Altern Med*. 2013 Sep 22;13:232.
- [2]. Quinn L, et al. Extraction and Quantification of Sinapinic Acid from Irish Rapeseed Meal and Assessment of Angiotensin-I Converting Enzyme (ACE-I) Inhibitory Activity. *J Agric Food Chem*. 2017 Aug 16;65(32):6886-6892.
- [3]. Zych M, et al. The Effects of Sinapic Acid on the Development of Metabolic Disorders Induced by Estrogen Deficiency in Rats. *Oxid Med Cell Longev*. 2018 Jun 4;2018:9274246.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA