Inhibitors

MCE MedChemExpress

D-Xylulose-1-¹³C

Molecular Weight:

Cat. No.: HY-W010256S

CAS No.: 131771-46-5 Molecular Formula: $C_4^{13}CH_{10}O_5$

Target: Endogenous Metabolite; Isotope-Labeled Compounds

Pathway: Metabolic Enzyme/Protease; Others

151.12

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.

O OH HQ_{3C} OH H₂ OH

BIOLOGICAL ACTIVITY

Description	D-Xylulose- 1^{-13} C is the 13 C labeled D-Xylulose. D-xylulose is a precursor of the pentiol D-arabi[1][2].
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

[2]. Huck JH, et al. Evaluation of pentitol metabolism in mammalian tissues provides new insight into disorders of human sugar metabolism. Mol Genet Metab. 2004 Jul;82(3):231-7.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA