

## Product Data Sheet

## Inhibitors • Screening Libraries • Proteins

## Uridine 5'-monophosphate-<sup>13</sup>C<sub>9</sub>,<sup>15</sup>N<sub>2</sub> disodium

Cat. No.:	HY-W013175S2	
Molecular Formula:	<sup>13</sup> C <sub>9</sub> H <sub>11</sub> <sup>15</sup> N <sub>2</sub> Na <sub>2</sub> O <sub>9</sub> P	H
Molecular Weight:	379.07	0-13C 0
Target:	Isotope-Labeled Compounds	$H^{13}C_{13}$
Pathway:	Others	$\begin{array}{c} H \\ H_{13}C \\ H_{13$
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	HO ISC H O H NaO

BIOLOGICAL ACTIVITY		
BIOLOGICALACIWIT		
Description	Uridine 5'-monophosphate- <sup>13</sup> C <sub>9</sub> , <sup>15</sup> N <sub>2</sub> (disodium) is the <sup>13</sup> C and <sup>15</sup> N labeled Uridine 5'-monophosphate disodium salt[1]. Uridine 5'-monophosphate disodium salt is component used for RNA synthesis[2].	
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

## REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

[2]. Hozumi Y, et al. Orotate phosphoribosyltransferase localizes to the Golgi complex and its expression levels affect the sensitivity to anti-cancer drug 5-fluorouracil. Biomed Res. 2015;36(6):403-9.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA