MCE MedChemExpress

Product Data Sheet

N-Acetyl-L-glutamic acid-d₅

Molecular Weight:

Target: Endogenous Metabolite; Endogenous Metabolite; Isotope-Labeled Compounds

Pathway: Metabolic Enzyme/Protease; Others

Storage: Powder -20°C 3 years

4°C 2 years
In solvent -80°C 6 months
-20°C 1 month

HO D D HN

BIOLOGICAL ACTIVITY

DescriptionN-Acetyl-L-glutamic acid. G₅ is the deuterium labeled N-Acetyl-L-glutamic acid. N-Acetyl-L-glutamic acid, a N-acyl-L-amino acid, is a component of animal cell culturing media. N-Acetyl-L-glutamic acid is a metabolite of Saccharomyces cerevisiae

and human[1].

In Vitro Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as

tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to

affect the pharmacokinetic and metabolic profiles of $drugs^{[1]}$.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

[2]. M Hashimoto, et al. Animal cell culturing media containing N-acetyl-L-glutamic acid. Patent.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA