


## N-Isovalerylglycine-d<sub>9</sub>

<b>Cat. No.:</b>	HY-W015464S		
<b>CAS No.:</b>	1330037-21-2		
<b>Molecular Formula:</b>	C <sub>7</sub> H <sub>4</sub> D <sub>9</sub> NO <sub>3</sub>		
<b>Molecular Weight:</b>	168.24		
<b>Target:</b>	Endogenous Metabolite		
<b>Pathway:</b>	Metabolic Enzyme/Protease		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### BIOLOGICAL ACTIVITY

<b>Description</b>	N-Isovalerylglycine-d <sub>9</sub> is the deuterium labeled N-Isovalerylglycine. N-Isovalerylglycine is an acyl glycine and could be used as a biomarker for the predisposition for weight gain and obesity.
<b>In Vitro</b>	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019;53(2):211-216.
- [2]. ARYL SULFONOHYDRAZIDES. US9500660B2.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA