


(R)-Azetidine-2-carboxylic acid

Cat. No.:	HY-W017755
CAS No.:	7729-30-8
Molecular Formula:	C ₄ H ₇ NO ₂
Molecular Weight:	101.1
Target:	ADC Linker; PROTAC Linkers
Pathway:	Antibody-drug Conjugate/ADC Related; PROTAC
Storage:	<div> <div>Powder</div> <div> -20°C 3 years 4°C 2 years </div> </div> <div> <div>In solvent</div> <div> -80°C 6 months -20°C 1 month </div> </div>



BIOLOGICAL ACTIVITY

Description	(R)-Azetidine-2-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). (R)-Azetidine-2-carboxylic acid is also a alkyl chain-based PROTAC linker that can be
IC ₅₀ & Target	Non-cleavable Linker
In Vitro	<p>ADCs are comprised of an antibody to which is attached an ADC cytotoxin through an ADC linker^[1].</p> <p>PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins^[2].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

REFERENCES

- [1]. Beck A, et al. Strategies and challenges for the next generation of antibody-drug conjugates. Nat Rev Drug Discov. 2017;16(5):315-337.
- [2]. Nalawansha DA, et al. PROTACs: An Emerging Therapeutic Modality in Precision Medicine. Cell Chem Biol. 2020;27(8):998-985.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA