


## Propionylpromazine-d<sub>6</sub> hydrochloride

<b>Cat. No.:</b>	HY-W040146S
<b>CAS No.:</b>	1262770-67-1
<b>Molecular Formula:</b>	C <sub>20</sub> H <sub>19</sub> D <sub>6</sub> ClN <sub>2</sub> OS
<b>Molecular Weight:</b>	382.98
<b>Target:</b>	Dopamine Receptor
<b>Pathway:</b>	GPCR/G Protein; Neuronal Signaling
<b>Storage:</b>	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



### BIOLOGICAL ACTIVITY

<b>Description</b>	Propionylpromazine-d <sub>6</sub> (hydrochloride) is the deuterium labeled Propionylpromazine hydrochloride. Propionylpromazine hydrochloride (Propiopromazine hydrochloride), a dopamine receptor D2 (DRD2) antagonist, can be used in the research of Parkinson disease[1].
<b>IC<sub>50</sub> &amp; Target</b>	D <sub>2</sub> Receptor
<b>In Vitro</b>	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019;53(2):211-216.
- [2]. Tyers, Mike, et al. Method for treating a brain cancer with ifenprodil. US8058243.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA