

Product Data Sheet

Fmoc-Asp-OMe

Cat. No.:HY-W041982CAS No.:145038-52-4Molecular Formula: $C_{20}H_{19}NO_6$ Molecular Weight:369.37

Target: Amino Acid Derivatives

Pathway: Others

Storage: Powder -20°C 3 years

4°C 2 years

In solvent -80°C 6 months

-20°C 1 month

SOLVENT & SOLUBILITY

In Vitro

DMSO: 100 mg/mL (270.73 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	2.7073 mL	13.5366 mL	27.0731 mL
	5 mM	0.5415 mL	2.7073 mL	5.4146 mL
	10 mM	0.2707 mL	1.3537 mL	2.7073 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.5 mg/mL (6.77 mM); Suspended solution; Need ultrasonic
- 2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.77 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	Fmoc-Asp-OMe is an aspartic acid derivative $^{[1]}$.
In Vitro	Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

1]. Luckose F, et al. Effects of a	mino acid derivatives on physi	cal, mental, and physiological	activities. Crit Rev Food Sci Nutr. 201	5;55(13):1793-1144.
	Courtiem, Dreaduct has not	hoon fully validated for me	odical applications. For years yet	use enly
	Tel: 609-228-6898	Fax: 609-228-5909	edical applications. For research E-mail: tech@MedChemExp	
			outh Junction, NJ 08852, USA	1000.00111

Page 2 of 2 www.MedChemExpress.com