## **Product** Data Sheet

## rU Phosphoramidite-<sup>13</sup>C<sub>2</sub>,d<sub>1</sub>

**Cat. No.:** HY-W048482S

Molecular Formula:  $C_{43}^{13}C_{2}H_{60}DN_{4}O_{9}PSi$ 


Molecular Weight: 864.04

Target: Isotope-Labeled Compounds; DNA/RNA Synthesis

Pathway: Others; Cell Cycle/DNA Damage

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.



## **BIOLOGICAL ACTIVITY**

Description	rU Phosphoramidite- $^{13}$ C <sub>2</sub> ,d <sub>1</sub> (DMT-2'O-TBDMS-rU phosphoramidite- $^{13}$ C <sub>2</sub> ,d <sub>1</sub> ) is deuterium and $^{13}$ C-labeled rU Phosphoramidite (HY-W048482). rU Phosphoramidite is a phosphorite monomer that can be used in the synthesis of oligonucleotides.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> .  MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## **REFERENCES**

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA