


## Butane-1,4-diylidiphosphonic acid

Cat. No.:	HY-W075674
CAS No.:	4671-77-6
Molecular Formula:	C <sub>4</sub> H <sub>12</sub> O <sub>6</sub> P <sub>2</sub>
Molecular Weight:	218.08
Target:	PROTAC Linkers
Pathway:	PROTAC
Storage:	<div>Powder</div> <div>-20°C 3 years</div> <div>4°C 2 years</div> <div>In solvent</div> <div>-80°C 6 months</div> <div>-20°C 1 month</div>



### BIOLOGICAL ACTIVITY

Description	Butane-1,4-diylidiphosphonic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs <sup>[1]</sup> .
IC <sub>50</sub> & Target	Alkyl-Chain
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA