## Sorbitan monooctadecanoate

Cat. No.:	HY-W12959	3	
CAS No.:	1338-41-6		
Target:	Biochemica	ıl Assay F	Reagents
Pathway:	Others		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month

SOLVENT & SOLUE	BILITY
In Vitro	Ethanol : 5 mg/mL (ultrasonic and warming and heat to 60°C)
In Vivo	<ol> <li>Add each solvent one by one: 10% EtOH &gt;&gt; 40% PEG300 &gt;&gt; 5% Tween-80 &gt;&gt; 45% saline Solubility: ≥ 0.5 mg/mL (Infinity mM); Clear solution</li> <li>Add each solvent one by one: 10% EtOH &gt;&gt; 90% corn oil Solubility: ≥ 0.5 mg/mL (Infinity mM); Clear solution</li> </ol>

## **BIOLOGICAL ACTIVITY**

Description	Sorbitan monooctadecanoate is a hydrophobic nonionic surfactant. Sorbitan monooctadecanoate can be used as an
	excipient, such as emulsifiers, nonionic surfactants, solubilizers, wetting and dispersing/suspending agents. Pharmaceutical
	excipients, or pharmaceutical auxiliaries, refer to other chemical substances used in the pharmaceutical process other than
	pharmaceutical ingredients. Pharmaceutical excipients generally refer to inactive ingredients in pharmaceutical
	preparations, which can improve the stability, solubility and processability of pharmaceutical preparations. Pharmaceutical
	excipients also affect the absorption, distribution, metabolism, and elimination (ADME) processes of co-administered drugs <sup>[1][2]</sup> .

## REFERENCES

[1]. Murdan S, et al. Novel sorbitan monostearate organogels[J]. Journal of pharmaceutical sciences, 1999, 88(6): 608-614.

[2]. Elder DP, et al. Pharmaceutical excipients - quality, regulatory and biopharmaceutical considerations. Eur J Pharm Sci. 2016 May 25;87:88-99.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA



**Product** Data Sheet