


## L-Tyrosine Hydrazide

<b>Cat. No.:</b>	HY-W142000
<b>CAS No.:</b>	7662-51-3
<b>Molecular Formula:</b>	C <sub>9</sub> H <sub>13</sub> N <sub>3</sub> O <sub>2</sub>
<b>Molecular Weight:</b>	195.22
<b>Target:</b>	Amino Acid Derivatives
<b>Pathway:</b>	Others
<b>Storage:</b>	4°C, protect from light, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light, stored under nitrogen)



### SOLVENT & SOLUBILITY

#### In Vitro

1M HCl : 50 mg/mL (256.12 mM; ultrasonic and adjust pH to 1 with HCl)  
DMSO : 5 mg/mL (25.61 mM; ultrasonic and warming and heat to 60°C)

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		5.1224 mL	25.6121 mL	51.2243 mL
	5 mM		1.0245 mL	5.1224 mL	10.2449 mL
	10 mM		0.5122 mL	2.5612 mL	5.1224 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

L-Tyrosine Hydrazide is a tyrosine derivative<sup>[1]</sup>.

#### In Vitro

Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances<sup>[1]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. Luckose F, et al. Effects of amino acid derivatives on physical, mental, and physiological activities. Crit Rev Food Sci Nutr. 2015;55(13):1793-1144.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA