


## Tetraglycine hydrochloride

<b>Cat. No.:</b>	HY-W142467A
<b>CAS No.:</b>	38126-71-5
<b>Molecular Formula:</b>	C <sub>8</sub> H <sub>15</sub> ClN <sub>4</sub> O <sub>5</sub>
<b>Molecular Weight:</b>	282.68
<b>Sequence Shortening:</b>	GGGG
<b>Target:</b>	Others
<b>Pathway:</b>	Others
<b>Storage:</b>	Sealed storage, away from moisture and light, under nitrogen
	Powder    -80°C    2 years
	-20°C    1 year



\* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light, under nitrogen)

### BIOLOGICAL ACTIVITY

<b>Description</b>	Tetraglycine hydrochloride is a oligopeptide composed of four glycine monomers <sup>[1]</sup> .								
<b>In Vivo</b>	<p>Tetraglycine and Triglycine (1.0 μmol glycine/g body wt (246.22 mg/kg), injected into a central vein) results in greater glycine concentration in the kidney than injection of either <a href="#">Glycine</a> (HY-Y0966) or Diglycine<sup>[1]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1"> <tr> <td><b>Animal Model:</b></td> <td>Male Sprague-Dawley rats (270-300 g)<sup>[1]</sup></td> </tr> <tr> <td><b>Dosage:</b></td> <td>1.0 μmol glycine/g body wt (246.22 mg/kg)</td> </tr> <tr> <td><b>Administration:</b></td> <td>IV, injected over a period of 30 s</td> </tr> <tr> <td><b>Result:</b></td> <td>Five minutes after the Tetraglycine injection, there were accumulations of diglycine, triglycine, and Tetraglycine in the kidney.</td> </tr> </table>	<b>Animal Model:</b>	Male Sprague-Dawley rats (270-300 g) <sup>[1]</sup>	<b>Dosage:</b>	1.0 μmol glycine/g body wt (246.22 mg/kg)	<b>Administration:</b>	IV, injected over a period of 30 s	<b>Result:</b>	Five minutes after the Tetraglycine injection, there were accumulations of diglycine, triglycine, and Tetraglycine in the kidney.
<b>Animal Model:</b>	Male Sprague-Dawley rats (270-300 g) <sup>[1]</sup>								
<b>Dosage:</b>	1.0 μmol glycine/g body wt (246.22 mg/kg)								
<b>Administration:</b>	IV, injected over a period of 30 s								
<b>Result:</b>	Five minutes after the Tetraglycine injection, there were accumulations of diglycine, triglycine, and Tetraglycine in the kidney.								

### REFERENCES

[1]. Adibi SA, et al. Enrichment of glycine pool in plasma and tissues by glycine, di-, tri-, and tetraglycine. Am J Physiol. 1982 Nov;243(5):E413-7.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA