


N-Acetyl-L-phenylalanine

Cat. No.:	HY-Y0068
CAS No.:	2018-61-3
Molecular Formula:	C ₁₁ H ₁₃ NO ₃
Molecular Weight:	207.23
Target:	Endogenous Metabolite
Pathway:	Metabolic Enzyme/Protease
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 100 mg/mL (482.56 mM; Need ultrasonic)
 H₂O : 7.69 mg/mL (37.11 mM; Need ultrasonic)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		4.8256 mL	24.1278 mL	48.2556 mL
	5 mM		0.9651 mL	4.8256 mL	9.6511 mL
	10 mM		0.4826 mL	2.4128 mL	4.8256 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: PBS
Solubility: 7.14 mg/mL (34.45 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (12.06 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (12.06 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (12.06 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	N-Acetyl-L-phenylalanine (N-Acetylphenylalanine), the principal acylamino acid in Escherichia coli, is synthesized from L-phenylalanine and acetyl-CoA ^[1] .
IC ₅₀ & Target	Human Endogenous Metabolite

CUSTOMER VALIDATION

- Laurea Magistrale in Biomedical Engineering, Politecnico di Milano. 2019 Jun.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Krishna RV, et, al. Enzymic synthesis of N-acetyl-L-phenylalanine in Escherichia coli K12. Biochem J. 1971 Oct;124(5):905-13.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA