## L-Cysteine-d<sub>3</sub>,<sup>15</sup>N

MedChemExpress

Cat. No.:	HY-Y0337S7		
CAS No.:	1795787-05-1		c
Molecular Formula:	C <sub>3</sub> H <sub>4</sub> D <sub>3</sub> <sup>15</sup> NO <sub>2</sub> S		
Molecular Weight:	125.17		
Target:	Endogenous Metabolite; Isotope-Labeled Compounds	HS´ 🏹 `OH	
Pathway:	Metabolic Enzyme/Protease; Others	$D^{15}NH_2$	
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	L	

Product Data Sheet

BIOLOGICAL ACTIVITY		
BIOLOGICAL ACTIVITY		
Description	L-Cysteine-d <sub>3</sub> , <sup>15</sup> N is the deuterium and <sup>15</sup> N-labeled L-Cysteine. L-Cysteine is a conditionally essential amino acid, which acts as a precursor for biologically active molecules such as hydrogen sulphide (H2S), glutathione and taurine. L-Cysteine suppresses ghrelin and reduces appetite in rodents and humans[1].	
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

## REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909

5909 E-mail: tech@MedChemExpress.com