

Product Data Sheet

HODHBt

Cat. No.:HY-Y0493CAS No.:28230-32-2Molecular Formula: $C_7H_5N_3O_2$ Molecular Weight:163Target:STAT; HIV

Talget.


Pathway: JAK/STAT Signaling; Stem Cell/Wnt; Anti-infection

Storage: Powder -20°C 3 years

In solvent

4°C 2 years -80°C 6 months

-20°C 1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO: 500 mg/mL (3067.48 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	6.1350 mL	30.6748 mL	61.3497 mL
	5 mM	1.2270 mL	6.1350 mL	12.2699 mL
	10 mM	0.6135 mL	3.0675 mL	6.1350 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

HODHBt (HOOBt) inhibits STAT5-SUMO interaction by blocking SUMOylation of phosphorylated STAT5. HODHBt enhances the magnitude of IL-15 signaling and significantly increases the natural killer (NK) cell cytotoxicity phenotype and function and the generation of cytokine-induced memory-like (CIML) natural killer (NK) cells. HODHBt can be used for research of

HIV-infection and cancer^[1].

IC₅₀ & Target STAT5

In Vitro HODHBt (100 μ M; 24 h, and 48 h respectively) increases IL-15-mediated STAT phosphorylation in NK cells and enhances the cytotoxic profile of NK cells^[1].

 $HODHBt~(100~\mu\text{M}; 24~h)~enhances~IL-15-mediated~(100~ng/mL)~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~of~NK~cells~against~HIV-infected~cells~(K562~cells)^{[1]}~cytotoxicity~cytotoxicity~of~NK~cells~against~hi~cytotoxicity~cytotoxic$

HODHBt (100 μ M; 24 h) enhances IL-15-mediated (100 ng/mL, 6 h) cytotoxicity of NK cells against cancer cell lines (A2780 and U877; OCILy1 and OCILy10)^[1].

HODHBt (100 μ M; 7 d) results in enhanced memory response upon recall in generation of human CIML NK cells in vitro^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES				
[1]. Macedo AB, et al. The HIV Latency Reversal Agent HODHBt Enhances NK Cell Effector and Memory-Like Functions by Increasing Interleukin-15-Mediated STAT Activation. J Virol. 2022 Aug 10;96(15):e0037222.				
	Caution: Product has not been fully validated for medical applications. For research use only.			
	Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com			
	Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA			

Page 2 of 2 www.MedChemExpress.com